

# Internal Jugular Vein Compression: A Novel Approach to Mitigate Blast Induced Hearing Injury

\*†Brian Sindelar, ‡Michael Shinnars, \*Sydney Sherman, §Kevin Novak, †Kristine Erickson  
\*Vimal Patel, †Paul Kubilis, ||David Smith, \*John Finan, and \*Julian E. Bailes

\*Department of Neurosurgery, NorthShore University HealthSystem, Evanston, Illinois; †Department of Neurosurgery, University of Florida, Gainesville, Florida; ‡Department of Surgery, Division of Otolaryngology; §Department of Neurophysiology, NorthShore University HealthSystem, Evanston, Illinois; and ||Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio

**Hypothesis:** Internal jugular vein (IJV) compression before blast injury will lead to reduced risk of traumatic hearing injury following exposure to a blast injury.

**Background:** IJV compression and its effects on not only intracranial, but also intracochlear pressure may potentiate blast induced hearing injury, therefore, precluding its use as a prophylactic therapy for blast induced traumatic brain injury.

**Methods:** Twenty Sprague Dawley rats were exposed to a  $17.9 \pm 0.4$  PSI (195.8 dB SPL) right sided shock wave in which 10 had application of a custom IJV compression collar before injury. All rodents received baseline and post blast injury otoacoustic emission (OAE) and auditory brainstem response (ABR) testing followed by cochlear histology.

**Results:** IJV compression was shown to significantly reduce ABR and OAE threshold shifts in comparison to the non-intervention group by:  $14.9 \pm 4.8$  dB (right ear ABR 0.5 kHz

Day 1 post blast,  $p=0.01$ ),  $13.1 \pm 4.9$  dB (right ear ABR 4 kHz Day 1 post blast,  $p=0.04$ ),  $16.5 \pm 4.5$  dB (right ear ABR click Day 1 post blast,  $p=0.003$ ),  $12.1 \pm 4.6$  dB (right ear ABR click Day 6 post blast,  $p=0.04$ ), and  $14.0 \pm 3.2$  dB (both ears OAE 3.2–10 kHz,  $p<0.0001$ ). Also, those animals with collar application had a greater number of total hair cells per mm from 70 to 100% distance from the cochlear apex following blast injury in comparison to those without intervention (blast:  $211.8 \pm 27.5$  versus blast+collar:  $355.5 \pm 39.5$  [ $p=0.0002$ ]).

**Conclusion:** This study supports the use of IJV compression in a pre-clinical model as a new prophylactic mechanism to combat blast induced hearing injury. **Key Words:** Blast—Internal jugular vein compression—Traumatic hearing injury.

*Otol Neurotol* 38:xxx-xxx, 2017.

Internal jugular vein (IJV) compression has been proposed as a prophylactic measure to diminish the microscopic and macroscopic injuries associated with traumatic brain injury (TBI). The preeminent theory is that a TBI is due to the differential acceleration/deceleration of the viscoelastic cerebrum within the skull leading to stretch and strain forces upon the cells, white matter, and blood vessels (1–6). These forces cause initial primary axotomy and hemorrhage along with an initiation of a complex neurometabolic and vascular response leading to alterations in neuronal functioning (7–13). Prophylactic mild IJV compression engorges the

cranial venous system, enlarges the brain, and increases brain turgor, therefore, reducing relative motion of the cerebrum within the skull (14). Preclinical studies in IJV compression before injury exposure have demonstrated its ability to reduce not only microscopic neuronal injury, but also the propensity for hemorrhagic extension (3,4,15). Recently, two clinical studies have successfully made the translational leap to demonstrate that use of an IJV compression collar in high school football and hockey players provided a significant reduction in sub-concussive axonal injury on diffusion tensor imaging following a season of play (16,17).

Application of this technology has been suggested for use in military personnel to combat military blast TBI, but concerns have been raised due to its potential consequences on blast induced hearing injury. Specifically, since IJV compression causes mild increases in intracranial volume and pressure and subsequently intracochlear pressure, what would be the effect of prophylactic IJV compression on traumatic hearing injury in those service members exposed to a blast injury (18–22)? With the already high prevalence of hearing deficits in military

Address correspondence and reprint requests to Julian Bailes, M.D., Department of Neurosurgery, Northshore University HealthSystem, Evanston Hospital, 3rd Floor Kellogg Bldg, Evanston, IL 60201; E-mail: jbailes@northshore.org

Funding: Funding for this research was provided by Q30 Sports Science, LLC.

JB holds stock in Q30 Sports Science, LLC based on the original patent for IJV compression, and DS is a consultant to Q30 Sports Science, LLC.

The authors disclose no conflicts of interest.

DOI: 10.1097/MAO.0000000000001332

service members, any therapeutic modality that could potentially increase prevalence or worsen its morbidity would not be clinically feasible (23).

This study attempts to address if IJV compression before blast injury will lead to increased or reduced risk of traumatic hearing injury following exposure to a blast injury, to evaluate the collar's potential as a means for TBI prophylaxis in the military. If suggestive of a therapeutic effect, outcomes of this study have the potential to revolutionize our approach to improving protection of our military personnel during operational theaters.

## METHODS

### Experimental Design Overview

To assess the effect of IJV compression on traumatic hearing injury, we exposed 20 rats to a right sided 18.0 PSI overpressure blast in which half of the animals had preblast application of a specialized IJV compression collar (blast+collar) and the other half did not (blast). All animals had baseline and post injury otoacoustic emissions (OAE) and auditory brainstem response (ABR) testing to assess functional injury of the cochlea and eighth cranial nerve, followed by structural evaluation via cochlear histology after euthanasia 9 days after injury.

### Animals

Twenty male adult (3–6 months old) Sprague Dawley rats (Charles River Laboratories, Wilmington, MA) weighing between 350 and 400 g were delivered 5 days before initial testing in order for acclimatization. The group was heterogeneous regarding inner ear anatomy like hair cell density, ear canal length etc., but maintaining comparable age and weight ranges along with randomization was assumed to provide for these differences. All animals were housed in the hospital animal facility with appropriate feed and water and maintained on normal night/day cycles. Animals were monitored daily by trained animal technicians by the oversight of the hospital veterinarian and all care, and the entirety of the experimental protocol was approved by the NorthShore Evanston Institutional Animal Care and Use Committee (EH15–391).

### Therapeutic IJV Compression Collar

To apply mild IJV compression, a customized collar, modeled from Smith et al. (4), was comprised of elastic with properly placed plastic beads to ensure compression of the bilateral IJV and avoid tracheal compromise or carotid stimulation. Before the clinical experiment, proper collar design (bead location and collar tightness) was determined by the trial of different collar prototypes in concert with invasive intracranial and non-invasive intraocular pressure monitoring. This also assured that proper collar application could be assessed experimentally through non-invasive measures without the need to alter the intracranial milieu through invasive monitoring.

All animals before blast injury received intraocular pressure (IOP) monitoring every 30 seconds for a total of 3 minutes. For those animals that were randomized to collar application before blast injury, the customized IJV compression collar was applied and IOP (Icare TONOLAB #TL-1, Colonial Medical Supply, Windham, NH) was again measured every 30 seconds for 3 minutes. The collar was re-adjusted and replaced for those animals that did not demonstrate immediate increases in intraocular pressure.

### Blast Induced Hearing Injury

For blast induced hearing injury, the animals were anesthetized with isoflurane (2–5% for induction and 1–3% for maintenance) along with one dose of buprenorphine (0.05 mg/kg) for analgesia. Once anesthetized, the animals received IOP recordings and collar application (if appropriate), followed by placement into a small steel cylinder surrounded by foam padding to protect the thorax and eliminate barotrauma. Attention was focused on maintaining a constant location for the head to be exposed, being at the craniocervical junction just distal to the ears. The animal was then positioned in an upright manner, perpendicular to the front of the blast tube opening to receive a right lateralized blast. All animals' heads were positioned an equal distance from the blast tube exit site, roughly 2 cm.

The shock tube apparatus used to recreate blast induced traumatic hearing injury has been validated in previous studies (24–26). To create a blast wave with this tube, compressed helium gas is pumped into the driver section that is sealed by mylar membrane (McMaster Carr, Elmhurst, IL). At a given pressure, based on the thickness of the mylar, the membrane will rupture sending a shock wave down the driven section, striking the animal that is situated at the open end of the chamber. Three piezoresistive pressure transducers (8530B-200, Endevco Meggitt, Irvine, CA) are situated at the end of the blast tube to record blast overpressure and duration. Verified before the experimental study, one 0.010" mylar membrane produces a 17 to 18 PSI blast overpressure with a duration of 0.4 ms. This level was chosen to reproduce a typical IED blast overpressure (between 7 and 145 PSI) and not to exceed the recommended rodent scaled blast duration of 0.3 to 3 ms; but, while also reducing traumatic tympanic membrane rupture (26–29). All animals had visual inspection of their tympanic membranes under light microscopy following blast exposure, and those with tympanic membrane rupture were removed from the final OAE analysis.

### Auditory Brainstem Responses

ABRs were performed the day previous (baseline), 1-day post, and 6-day post blast injury using Nihon Kohden (Irvine, CA) MEE clinical evoked potential system. All animals were anesthetized with inhaled isoflurane, placed on a warming pad within a sound insulating container, and received continuous pulse oximetry and heart rate monitoring for the duration of testing. Following successful sedation, three subdermal scalp recording electrodes were placed in sterile fashion at the vertex and at both ears. Sound click and toneburst stimuli were delivered with foam inserts fitted tightly in the ear canal. Broadband (range of 3–8 kHz range, peak of 3–4 kHz) rarefaction clicks of 0.1 ms duration were delivered at 27.1 stimuli per second, and scalp responses (Cz-Ai) averaged over 200 to 2000 stimuli, depending on the signal to noise ratio. Similarly, two frequencies of toneburst stimuli were delivered at 27.1 stimuli/s: 500 Hz burst of 4 ms duration (2 ms rise, 0 ms plateau, 2 ms decay), and 4000 Hz wave of 6 ms duration (3 ms rise, 0 ms plateau, 3 ms decay). Contralateral masking noise was used for all ABRs. ABR thresholds were determined visually for 500 Hz toneburst, 4000 Hz toneburst, and broadband clicks for each ear by following Wave II response down from 80 dB and noting the intensity when it disappeared. ABR threshold shifts were calculated by subtracting post exposure from baseline testing values.

### Distortion Product Otoacoustic Emissions

OAEs were performed the day prior (baseline), 5-minutes post, 1-day post, and 6-day post blast injury. Animals were

again prepared in similar fashion as the ABR testing. Following sedation, single ear rubber tipped ear inserts were placed, assuring a tight fit within the ear canal. Through the use of the Interacoustics Titan DPOAE440 (e3 Gordon Stowe, Birmingham, AL), cochlear emissions were measured from 2 to 10 kHz using two primary tones,  $f_1$  and  $f_2$  levels of  $L_1$  of  $L_2 = 65$  of 55 dB with a ratio  $f_2$  of  $f_1 = 1.22$ . OAE levels were calculated by subtracting pre and post OAE levels. Also, the noise floor level was used for calculating OAE level changes in those that fell below the noise level floor for a specific frequency.

### Histological Examination

Following completion of testing, the rats were euthanized by decapitation under deep anesthesia 9 days following blast exposure. Twelve total animals (six blast+collar and six blast) were randomly selected for cochlear histology. The temporal bones were quickly removed and fixed in 10% neutral buffered formalin for 2 to 5 days. The cochlear basilar membrane was dissected out under a dissection microscope, and stained with Harris' hematoxylin solution for 5 minutes. To remove the undesirable coloration, specimens were differentiated with 0.3% acid alcohol for few seconds, and then the samples were rinsed with running tap water to convert the color into violet blue. The cochlear basilar membrane was further trimmed and mounted in glycerin on glass slides as a flat surface preparation. The stereocilia, cuticular plate, and nucleus of hair cells were identified under a Zeiss microscope at 400 times magnification. The number of inner and outer hair cells was counted over successive 0.24 mm segments along the entire length of the cochlear basilar membrane. Hair cell density was computed into 10% segments of the cochlea from apex to base, and these values were normalized to lab standards based on hair cell counts from six young, healthy rats. A cochleogram was constructed by plotting the percent inner and outer hair cell loss as a function of the percent distance from the apex of the cochlea. For analysis, individual cochleograms were grouped to generate a mean cochleogram for each experimental condition using the custom software.

### Statistical Analysis

A linear mixed effect model was used to assess the effects of experimental factors on both ABR and OAE threshold level shifts relative to baseline levels. Treatment (blast+collar versus blast) was modeled as a between-subject fixed effect; ear (right/left), test day, and ABR/OAE test frequency were modeled as within-subject fixed effects; and rat was modeled as a subject random effect. Type III tests of fixed effects and sub-effects in the model were evaluated using  $F$  tests. Collar minus control differences in response were evaluated using  $t$  statistics, with adjustment of  $p$  values and confidence intervals within logical groupings of tests using Sidak's method to maintain group wise significance levels of 0.05.

A generalized linear mixed effect models with Poisson errors (IHC hair counts) or Negative Binomial errors (OHC hair counts) were used to assess the effects of experimental factors on cochlear hair counts. Treatment (blast+collar versus blast) was modeled as a between-subject fixed effect; ear (right/left), and distance along the cochlea were modeled as within-subject fixed effects; and rat was modeled as a subject random effect. Average distance-specific hair counts from six healthy control animals were used as offsets in each model, allowing experimental factor means to be interpreted as the proportion of cells counted relative to "ideal" hair counts from typical healthy

rats. Type III tests of fixed effects and sub-effects in the model were evaluated using  $F$  tests. Collar versus control ratios of mean standardized responses (hair count proportions) were evaluated using  $t$  statistics, with adjustment of  $p$  values and confidence intervals within logical groupings of tests using Sidak's method to maintain groupwise significance levels of 0.05.

Lastly, blast pressure, blast duration, and IOP are expressed as mean  $\pm$  standard deviation while ABR, OAE, and cochlear histology is expressed as mean  $\pm$  standard error of the mean.

## RESULTS

### Blast Exposure $\pm$ Collar

Following anesthesia, all animals received IOP pressure monitoring for 3 minutes to determine a baseline IOP of  $13.1 \pm 2.5$  mm Hg. After application of our customized IJV compression collar, the IOP increased to  $24.3 \pm 1.8$  mm Hg. All animals tolerated application of the IJV compression collar without any alterations in monitored pulse oximetry measurements.


The 20 animals were then exposed to a  $17.9 \pm 0.4$  PSI (195.8 dB SPL) blast overpressure with a duration of  $0.4 \pm 0.03$  ms followed by immediate removal of the IJV collar. There was no statistical difference in blast wave mechanics between the blast+collar and blast group (blast amplitude or duration). With this blast over pressure, none of the animals had a ruptured tympanic membrane in the left ear, but one (blast+collar) and three (blast) animals had a right ear tympanic membrane rupture (not statistically significant). These animals were removed from the final OAE analysis.

### ABR Threshold Shifts

The left ear (farthest from the blast wave), showed no statistically significant difference between the blast+collar and blast animals. Interestingly, all ABR stimuli (500, 4000 Hz, and click) presented to the right ear showed a statistically significant improvement in threshold shifts in those animals with blast+collar 1 day following blast injury. In more detail, collar application reduced the ABR threshold shift by  $14.9 \pm 4.8$  dB (0.5 kHz,  $p = 0.01$ ),  $13.1 \pm 4.9$  dB (4 kHz,  $p = 0.04$ ), and  $16.5 \pm 4.5$  dB (click,  $p = 0.003$ ). Six days following the blast injury, threshold level shifts of 500, 4000 Hz, and click demonstrated partial recovery in the blast animals and almost complete return to baseline in the blast+collar animals. Six days following injury, the click ABR stimuli still demonstrated a statistically significant reduction in threshold shift between the blast and blast+collar animals by  $12.1 \pm 4.6$  ( $p < 0.05$ ), but the 500 and 4000 Hz lost significance. The persistent difference seen in the click stimuli is likely due to the broadband frequency and its ability to assess a wider range of hair cells compensating for the improved temporary threshold shifts.

### OAE Threshold Shifts

A graphical analysis demonstrating the baseline OAEs followed by OAEs obtained 5 minutes, 1 day, and 6 days



**FIG. 1.** OAE responses at baseline (pre-blast), 5 minutes (Day 0), 1 day, and 6 days post blast in the right ear (blast side). *A*, Blast ( $n=7$ ) and *B*, blast+collar ( $n=9$ ). OAE indicates otoacoustic emission.

post blast injury in both right and left ears of the blast and blast+collar animals is depicted in Figures 1 and 2. Of note, the Sprague Dawley rats used for this experiment were shown to not have significant OAE responses below 3.2 kHz. There was a non-statistically significant trend of OAE threshold shifts to worsen from the assessment immediately following the blast injury to Day 1, but then appeared to improve at Day 6. This observation in threshold shift has been previously published (30). Figure 3 shows the total OAE threshold level shifts from baseline to Day 6 in both the right and left ears of the blast and blast+collar animals. Statistical significance is depicted on the graphs as  $* < 0.05$ ,  $** < 0.01$ , and  $*** < 0.001$ . Averaged over 3.2 to 10 kHz (the frequencies with substantial OAE responses at baseline in this cohort of Sprague Dawley rats), there was a statistically significant reduction in OAE threshold shifts in the blast+collar compared with the blast groups by  $12.5 \pm 3.3$  dB ( $p < 0.005$ ) in the left ear,  $15.5 \pm 3.3$  dB ( $p < 0.0001$ ) in the right ear, or  $14.0 \pm 3.2$  ( $p < 0.0001$ ) in both ears favoring the blast+collar group.


*Otology & Neurotology, Vol. xx, No. xx, 2017*

## Histology



Following euthanasia at 9 days post blast injury, the cochleograms demonstrated a significant reduction in inner (IHC) and outer (OHC) hair cell loss specifically at the location closest to the oval window of the cochlea, corresponding to higher frequencies (roughly 32 kHz).

For the left ear, the blast+collar had a significant reduction in IHC loss compared with the blast animals located at a distance of 90 to 100% from the apex (blast 62% versus blast+collar 10%,  $p < 0.0001$ ). Statistical significance was also seen at 80 to 90% and 90 to 100% distance from the apex in outer hair cell loss between the two groups (80 to 90% cochlear distance: blast 78% versus blast+collar 20%,  $p < 0.0001$ , 90 to 100% cochlear distance: blast 100% versus 39%,  $p < 0.05$ ) (Fig. 4).

For the right ear, the blast animals had a significant higher percentage of IHC loss located between 90 and 100% of the distance from the apex (blast 61% versus blast+collar 27%,  $p < 0.0001$ ). Statistical significance was also seen at 80 to 90% and 90 to 100% distance from the apex in outer hair cell loss between the two



**FIG. 2.** OAE responses at baseline (pre-blast), 5 minutes (Day 0), 1 day, and 6 days post blast in the left ear (opposite blast side). *A*, Blast ( $n=10$ ) and *B*, blast+collar ( $n=10$ ). OAE indicates otoacoustic emission.



**FIG. 3.** OAE threshold shifts from baseline to Day 6 post blast in the *A*, right ear (blast side) and *B*, left ear (opposite blast side). \* $p < 0.05$  versus blast, \*\* $p < 0.01$  versus blast, \*\*\* $p < 0.001$  versus blast. OAE indicates otoacoustic emission.



groups (80–90% cochlear distance: blast 77% versus blast+collar 44%,  $p < 0.0001$ , 90–100% cochlear distance: blast 100% versus 62%,  $p < 0.05$ ) (Figs. 4 and 5).

If specifically analyzing the 70 to 100% distance from the apex in the blast+collar compared with the blast group in all ears, there is a greater number of total hair cells (IHC+OHC) per mm in the blast+collar versus blast group following exposure (blast:  $211.8 \pm 27.5$  versus blast+collar:  $355.5 \pm 39.5$  [ $p = 0.0002$ ]). In comparison to six control animals, the blast+collar group did not have a statistically different total number of hair cells per mm at the 70 to 100% distance (blast+collar:  $355.5 \pm 39.5$  versus control:  $463.72 \pm 5.1$  [ $p > 0.05$ ]) (Fig. 6).


**DISCUSSION**

IJV compression has been shown in preclinical and clinical studies to have a potential benefit in mitigating

the injury that occurs in TBI, and therefore has further been proposed for use in military blast TBI. This study specifically attempted to determine if prophylactic IJV compression, and resultant changes in intracranial and, potentially, intracochlear pressure, would improve or worsen traumatic blast audiological injury. In this study, it was demonstrated that IJV compression before injury did not increase the functional thresholds (ABR and OAE) or exacerbate structural (cochlear histology) inner ear pathology following a blast injury. Interestingly, this novel therapy was actually shown to dramatically reduce traumatic hearing injury in the rodent model based on ABR, OAE, and cochlear histology. Though difficult to compare to Li et al.'s (31) study assessing hearing injury in guinea pigs with ear inserts exposed to a shock wave due to differing methods (blast PSI, animal and blast model, time frame ABR obtained post blast, and



**FIG. 4.** Percent outer hair cell (OHC) and inner hair cell (IHC) loss compared with normal control graphed as a percent distance from the apex of the cochlea in the blast+collar and blast animals following euthanasia 9 days post blast injury. *A*, Right ear (blast side) and *B*, left ear (opposite blast side).



**FIG. 5.** Typical cochlear histology obtained from A, blast and B, blast+collar groups. Note the dramatic IHC and OHC loss seen in the blast animal. Images shown are from right ear (blast side) in the region 60 to 80% from apex. IHC indicates inner hair cell; OHC, outer hair cell.


euthanasia dates for histology), the similar and in some cases inferior outcomes of ear inserts compared with IJV compression has the potential to completely revolutionize our approach to prophylactic mitigation of blast traumatic hearing injury (31).

Communication from the intracranial to the intracochlear space has been demonstrated in animal models likely through a widely patent cochlear aqueduct (18–22,32–35). Presumably, IJV compression resulting in mild increases in intracranial volume and pressure could be transmitted to the cochlea via this route. The resultant change in intracochlear dynamics potentially may offer either mechanical impedance against the acoustic or shock wave, or may alter the stiffness of the sensory epithelium resulting in reduced movement and tearing of hair cells as the wave travels through the cochlea (36–38). Though inspection of tympanic membranes was performed to assess for gross rupture, it is not possible to exclude micro-perforation. Therefore, postulating whether hair cell injury occurs due to the blast shock wave (rupturing the tympanic membrane) or acoustic sound wave (causing vibration of the tympanic membrane) traveling through the cochlea is not able to be determined. But, it is interesting that our change to the cochlear histology was significant for a reduction in hair cell loss occurring closest to the oval window where the shock wave would first enter the cochlea and cause substantial injury.

Though this research has the potential to dramatically alter our approach to hearing protection, the translatability of these findings from preclinical to clinical subjects is uncertain due to the possible differences in temporal bone anatomy. An anatomical dissection of human temporal bones performed by Allen et al. (39) demonstrated that only 34% had a patent cochlear aqueduct, 59% had a lumen which was filled with connective tissue,

and only 7% had either complete obliteration or occlusion by bone. It is unknown to what extent intracranial pressure modifies intracochlear pressure specifically in those patients with connective tissue filling the lumen. Therefore, the viability of IJV compression via this mechanism could potentially only benefit 34%, but may also benefit 93% of humans. Though the patency of the cochlear aqueduct in humans is questioned (20,38–45), multiple clinical studies have demonstrated alterations in inner ear pathology (OAE threshold changes and tympanic membrane displacement) in patients with elevated intracranial pressure (46–49). Due to these positive findings, non-invasive acoustic tests have been suggested for use as an indirect measure of intracranial pressure in patients suffering from traumatic brain injury or hydrocephalus (21,46–49). Until further research is done, the conduit (cochlear aqueduct, cochlear vein, perineural space, endolymphatic sac, etc.) between the intracranial and intracochlear space in humans can only be speculated (20,21,39,50). But, this discussion is futile because a connection does appear to exist, whatever that may be, to transmit ICP to the cochlea, making the hypothesis that IJV compression would also be feasible in clinical studies of traumatic hearing loss mitigation.

Aside from the preclinical nature of our rodent model, other limitations to this study include the lack of scientific inquiry into the mechanism of IJV compression, cochlear physiology, and its effects on shock and acoustic waves. For this reason, we can only provide speculation for these. Secondly, our functional and structural findings do not correlate with each other. Specifically, the statistically significant differences in OAE findings from roughly 5 to 10 kHz in the right and left ears did not show any significant pathology on cochlear histology. Due to the preliminary nature of this data, an early euthanasia time point was chosen. For this reason, lack of corresponding cochlear histology may be due to euthanasia that proceeded completion of hair cell death. Alternatively, sufficient time may have elapsed for hair cell atrophy and sloughing, but these findings could be suggestive of a recoverable functional injury occurring



**FIG. 6.** Total hair cell count per mm at a distance of 70 to 100% from the cochlear apex in the blast+collar (n = 12), blast (n = 12), and control (n = 6). \* $p < 0.01$  versus blast.

at 5 to 10 kHz, i.e., temporary threshold shift. Follow-up studies will require acute and chronic time points to evaluate these questions.

## CONCLUSION

IJV compression before blast induced hearing injury was shown to reduce both functional and structural measures of inner ear pathology in a rodent model (ABR, OAE, and cochlear histology). This fascinating study proposes a new prophylactic mechanism that not only provides benefits to TBI mitigation, but also may revolutionize our approach to traumatic hearing injury in both the military and possibly even civilian population.

## REFERENCES

- Ommaya AK, Goldsmith W, Thibault L. Biomechanics and neuropathology of adult and paediatric head injury. *Br J Neurosurg* 2002;16:220–42.
- Meaney DF, Morrison B, Dale Bass C. The mechanics of traumatic brain injury: a review of what we know and what we need to know for reducing its societal burden. *J Biomech Eng* 2014;136:021008.
- Turner RC, Naser ZJ, Bailes JE, Smith DW, Fisher JA, Rosen CL. Effect of slosh mitigation on histologic markers of traumatic brain injury: laboratory investigation. *J Neurosurg* 2012;117:1110–8.
- Smith DW, Bailes JE, Fisher JA, Robles J, Turner RC, Mills JD. Internal jugular vein compression mitigates traumatic axonal injury in a rat model by reducing the intracranial slosh effect. *Neurosurgery* 2012;70:740–6.
- Stone JL, Patel V, Bailes JE. The history of neurosurgical treatment of sports concussion. *Neurosurgery* 2014;75:S3–23.
- Bayly PV, Cohen TS, Leister EP, Ajo D, Leuthardt EC, Genin GM. Deformation of the human brain induced by mild acceleration. *J Neurotrauma* 2005;22:845–56.
- Guskiewicz KM, Mihalik JP. Biomechanics of sport concussion: quest for the elusive injury threshold. *Exerc Sport Sci Rev* 2011;39:4–11.
- McKee AC, Stein TD, Kiernan PT, Alvarez VE. The neuropathology of chronic traumatic encephalopathy. *Brain Pathol* 2015;25:350–64.
- Choe MC, Giza CC. Diagnosis and management of acute concussion. *Semin Neurol* 2015;35:29–41.
- Caskey RC, Nance ML. Management of pediatric mild traumatic brain injury. *Adv Pediatr* 2014;61:271–86.
- Hovda DA. The neurophysiology of concussion. *Prog Neurol Surg* 2014;28:28–37.
- Giza CC, Hovda DA. The neurometabolic cascade of concussion. *J Athl Train* 2001;36:228–35.
- Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery* 2014;75:S24–33.
- Leach J, Smith D, Myer G. Mild neck compression alters intracranial venous sinus volume: implications for a novel neuroprotective effect in concussion. Paper presented at: American Society of Neuroradiology Annual Meeting; 2013.
- Sindelar B, Bailes JE, Sherman SA, et al. Effect of internal jugular vein compression on intracranial hemorrhage in a porcine controlled cortical impact model. *J Neurotrauma* 2016. [Epub ahead of print]. doi:10.1089/neu.2016.4648.
- Myer GD, Yuan W, Barber Foss KD, et al. The effects of external jugular compression applied during head impact exposure on longitudinal changes in brain neuroanatomical and neurophysiological biomarkers: a preliminary investigation. *Front Neuro* 2016;7:74.
- Myer GD, Yuan W, Barber Foss KD, et al. Analysis of head impact exposure and brain microstructure response in a season-long application of a jugular vein compression Collar: a prospective, neuroimaging investigation in American football. *Br J Sports Med* 2016. [Epub ahead of print]. doi: 10.1136/bjsports-2016-096134.
- Suzuki T, Nakashima T, Yanagita N. Effects of increased cerebrospinal fluid pressure on cochlear and cerebral blood flow. *Eur Arch Otorhinolaryngol* 1993;250:332–6.
- Allen GW, Habibi M. The effect of increasing the cerebrospinal fluid pressure upon the cochlear microphonics. *Laryngoscope* 1962;72:423–34.
- Bachor E, Byahatti S, Karmody CS. New aspects in the histopathology of the cochlear aqueduct in children. *Am J Otol* 1999;20:612–20.
- Carlborg BI, Farmer JC Jr. Transmission of cerebrospinal fluid pressure via the cochlear aqueduct and endolymphatic sac. *Am J Otol* 1983;4:273–82.
- Sakka L, Chomicki A, Gabrillargues J, Khalil T, Chazal J, Avan P. Validation of a noninvasive test routinely used in otology for the diagnosis of cerebrospinal fluid shunt malfunction in patients with normal pressure hydrocephalus. *J Neurosurg* 2016;124:342–9.
- Gondusky JS, Reiter MP. Protecting military convoys in Iraq: an examination of battle injuries sustained by a mechanized battalion during Operation Iraqi Freedom II. *Mil Med* 2005;170:546–9.
- Effgen GB, Hue CD, Vogel E 3rd, et al. A Multiscale approach to blast neurotrauma modeling: Part II: methodology for inducing blast injury to in vitro models. *Front Neurol* 2012;3:23.
- Gullotti DM, Beamer M, Panzer MB, et al. Significant head accelerations can influence immediate neurological impairments in a murine model of blast-induced traumatic brain injury. *J Biomech Eng* 2014;136:091004.
- Panzer MB, Matthews KA, Yu AW, Morrison B 3rd, Meaney DF, Bass CR. A multiscale approach to blast neurotrauma modeling: Part I—development of novel test devices for in vivo and in vitro blast injury models. *Front Neurol* 2012;3:46.
- Mediavilla Varas J, Philippens M, Meijer SR, et al. Physics of IED blast shock tube simulations for mTBI research. *Front Neurol* 2011;2:58.
- Panzer MB, Wood GW, Bass CR. Scaling in neurotrauma: how do we apply animal experiments to people? *Exp Neurol* 2014;261:120–6.
- Needham CE, Ritzel D, Rule GT, Wiri S, Young L. Blast testing issues and TBI: experimental models that lead to wrong conclusions. *Front Neurol* 2015;6:72.
- Patterson JH Jr, Hamernik RP. Blast overpressure induced structural and functional changes in the auditory system. *Toxicology* 1997;121:29–40.
- Li CJ, Zhu PF, Liu ZH, et al. Comparative observation of protective effects of earplug and barrel on auditory organs of guinea pigs exposed to experimental blast underpressure. *Chin J Traumatol* 2006;9:242–5.
- Caye-Thomassen P, Worsøe L, Brandt CT, et al. Routes, dynamics, and correlates of cochlear inflammation in terminal and recovering experimental meningitis. *Laryngoscope* 2009;119:1560–70.
- Salt AN, Kellner C, Hale S. Contamination of perilymph sampled from the basal cochlear turn with cerebrospinal fluid. *Hear Res* 2003;182:24–33.
- Sziklai I, Ferrary E, Sterkers O, Amiel C. Vasopressin entry into the inner ear fluids of the rat. *Hear Res* 1987;29:245–50.
- Berndt H, Bergmann K, Wagner H, Gerhardt HJ. [The effects of methodical factors on the microphonic potential in the guinea pig cochlea with perfusion of perilymph]. *Acta Otolaryngol* 1975;79:372–7.
- Baird J. Explosive shocks and impedance mismatch in armatures. *Electromagn Phenom* 2003;3:405–13. Available at: <http://www.emph.com.ua/11/pdf/bair.pdf>. Accessed September 9, 2016.
- Nakagawa A, Manley GT, Gean AD, et al. Mechanisms of primary blast-induced traumatic brain injury: insights from shock-wave research. *J Neurotrauma* 2011;28:1101–19.
- Gopen Q, Rosowski JJ, Merchant SN. Anatomy of the normal human cochlear aqueduct with functional implications. *Hear Res* 1997;107:9–22.
- Allen GW, Dallos P, Sakamoto S, Homma T. Cochlear microphonic potential in cats. Effects of perilymphatic pressure. *Arch Otolaryngol* 1971;93:388–96.
- Yilmazer C, Sennaroglu L, Basaran F, Sennaroglu G. Relationship of the cochlear aqueduct and inner ear pressure in Ménière's disease and in a normal population. *Otol Neurotol* 2001;22:534–8.

41. Hongjian L, Guangke W, Song M, Xiaoli D, Daoxing Z. The prediction of CSF gusher in cochlear implants with inner ear abnormality. *Acta Otolaryngol* 2012;132:1271–4.
42. Ding X, Tian H, Wang W, Zhang D. Cochlear implantation in China: review of 1,237 cases with an emphasis on complications. *ORL J Otorhinolaryngol Relat Spec* 2009;71:192–5.
43. Buchman CA, Copeland BJ, Yu KK, Brown CJ, Carrasco VN, Pillsbury HC 3rd. Cochlear implantation in children with congenital inner ear malformations. *Laryngoscope* 2004;114:309–16.
44. Kim BG, Sim NS, Kim SH, Kim UK, Kim S, Choi JY. Enlarged cochlear aqueducts: a potential route for CSF gushers in patients with enlarged vestibular aqueducts. *Otol Neurotol* 2013;34:1660–5.
45. Schuknecht HF, Reisser C. The morphologic basis for perilymphatic gushers and ooze. *Adv Otorhinolaryngol* 1988;39:1–12.
46. Gwer S, Sheward V, Birch A, et al. The tympanic membrane displacement analyser for monitoring intracranial pressure in children. *Childs Nerv Syst* 2013;29:927–33.
47. Reid A, Marchbanks RJ, Burge DM, et al. The relationship between intracranial pressure and tympanic membrane displacement. *Br J Audiol* 1990;24:123–9.
48. Samuel M, Burge DM, Marchbanks RJ. Tympanic membrane displacement testing in regular assessment of intracranial pressure in eight children with shunted hydrocephalus. *J Neurosurg* 1998;88:983–95.
49. Wunderlich K, Brinker T, Ernst A. The correlation of tympanic membrane displacement measurements to direct intracranial pressure recordings in cat. In: Ernst A, Marchbanks R, editors. *Intracranial and Intralabyrinthine Fluids: Basic Aspect and Clinical Applications*. Berlin, Germany: Springer; 1996.
50. Lo WW, Daniels DL, Chakeres DW, et al. The endolymphatic duct and sac. *AJNR Am J Neuroradiol* 1997;18:881–7.